arXiv:1304.3825v1 [physics.class-ph] 13 Apr 2013

Maxwell stress analysis of displacement induced electric force and its application to natural self-oscillation of a free electron

Zhixian Yu
Department of Physics and Astronomy, University of New Mexico, Albuquerque NM 87131 USA and
College of Physics Science, Qingdao University, Qingdao 266071 China

Liang Yu*
Laboratory of Physics, Jiaming Energy Research, Qingdao 266003 China

(Dated: April 16, 2013)

The Maxwell stress analysis is used for the calculation of displacement induced temporary electric force of a single charge. This force comes from the variation of the charge's electric intensities that follow Coulomb's inverse square law, and is a kind of displacement dependent temporary restore force. The possibility of natural self-oscillation of a free electron which is driven by this self-force is also suggested.

PACS numbers: 03.50.De

INTRODUCTION

The distance dependence of the electrostatic force of Coulomb's inverse square law has been tested in high precision through different methods, see for example the discussion of Jackson[1] and experiment of Williams et al[2].

In this paper we want to show that some temporary displacement dependent electric force may be generated in relation to the Coulomb's law. This is a kind of self electric force. It is induced by any displacement of the electric charge which changes the distribution of its electric fields that follow Coulomb's law.

THE VARIATION OF DISTRIBUTION AND MAGNITUDE OF ELECTRIC INTENSITY OF A CHARGE SUBJECT TO TEMPORARY DISPLACEMENT

For a particle with electric charge e at rest at origin O first, according to Coulomb's law the distribution of its spherically symmetric electric intensity E follows the inverse square law, that is

$$E = \frac{e}{r_e^2}, \tag{1}$$

where r_e is magnitude of position vector from the particle to a field point. Suppose this charge is shifted along z axis by a displacement Z_e in a time interval $\triangle t$ and become stationary again at a position O_e. Thus the electromagnetic fields depending on its velocity and acceleration may be neglected first. By this displacement Z_e, the electric intensity E of Eq. (1) will change in both magnitude and distribution. During this time interval any variation signal of the electromagnetic fields will propagate first from the origin O to a spherical shell with radius $c\Delta t$ as shown in Fig. 1, where c is velocity of light. At a position P on this spherical shell, the distance to origin O is the radius $r = c\Delta t$ and to the new origin O_e is $[r_e]$. The relation between $[r_e]$ and $r = c\Delta t$ is that

$$[r_e]^2 = (c\Delta t)^2 + Z_e^2 - 2c\Delta t Z_e \cos\theta, \tag{2}$$

where θ is the angle between r and z axis. Now the electric field outside this spherical shell does not change, the electric intensity on the outer boundary of this surface at P is

$$E_{out} = \frac{e}{r^2} = \frac{e}{(c\Delta t)^2}. \tag{3}$$

The electric intensity on the inner boundary of this surface at P is

$$E_{in} = \frac{e}{[r_e]^2}. \tag{4}$$

Since by Eq. (2) $[r_e]^2 \neq (c\Delta t)^2$, these two electric intensities of Eq. (3) and Eq. (4) are not equal. But according to the requirement of div$\mathbf{E} = 0$ in charge free space, they should be equal in magnitude and direction. Although this requirement is applied for the total electric field, it is also correct for the Coulomb field alone here. Since E_{out} of Eq. (3) is the primary value, it must remain unchanged, thus E_{in} should be modified to fulfil the requirement of div$\mathbf{E} = 0$, that is to change E_{in} to

$$E_{in} = \frac{e}{[r_e]^2} \cdot \frac{[r_e]^2}{(c\Delta t)^2} = \frac{e}{{r_e}^2} MF = \frac{e}{(c\Delta t)^2}, \tag{5}$$

where MF is a kind of modification function. From Eq. (5) it is defined as

$$MF = \frac{[r_e]^2}{(c\Delta t)^2} = 1 + \frac{Z_e^2}{(c\Delta t)^2} - \frac{2Z_e \cos\theta}{c\Delta t}. \tag{6}$$

This kind of modification may also be seen from the relation of the solid angles $d\Omega$ and $d\Omega_e$ subtending to an area element ds at position P from the origins at O and



FIG. 1. The relations between $r = c\Delta t$ and $[r_e]$.

O_e respectively as shown in Fig. 2. For the two solid angles we have

$$d\Omega = \frac{ds}{r^2} = \frac{ds}{(c\Delta t)^2}, \tag{7}$$

$$d\Omega_e = \frac{ds}{[r_e]^2}, \tag{8}$$

and following Eq. (6) we have

$$\frac{d\Omega}{d\Omega_e} = \frac{[r_e]^2}{r^2} = \frac{[r_e]^2}{(c\Delta t)^2} = MF. \tag{9}$$

Since the number of electric lines within solid angle $d\Omega_e$ is the same as that within $d\Omega$, the area density of electric lines which represents the electric intensity will change by the ratio MF. This modification now not only modifies the electric intensity at position P, but should also be down to the whole solid angle $d\Omega_e$. Thus we may write the electric intensity within the solid angle $d\Omega_e$ as

$$E_e = \frac{e}{r_e^2} MF. \tag{10}$$

Then when the charge is subject to a temporary displacement Z_e during time interval Δt, the electric intensities should be changed totally according to Eq. (10) within the spherical surface of radius $r = c\Delta t$.

MAXWELL STRESS ANALYSIS OF THE DISPLACEMENT DEPENDENT ELECTRIC FORCE ON A CHARGE SUBJECT TO A TEMPORARY DISPLACEMENT



FIG. 2. The relation between solid angles $d\Omega_e$ and $d\Omega$.

It is known that electromagnetic force on a system may be calculated from the Maxwell stress on the surface that encloses this system. We see from above that for a particle with charge e and subject to a temporary displacement Z_e, the distribution and magnitude of its electric intensity follows the modified Coulomb's law as Eq. (10). Now we want to show that this particle will get a displacement dependent electric force generated by its own electric field, and this can be calculated from the Maxwell stress surrounding this particle. Now we may take a spherical surface with center at O_e and radius a surrounding this particle, and according to Eq. (10), the electric intensity at this surface E_a as shown in Fig. 3 is

$$E_a = \frac{e}{a^2} MF. \tag{11}$$

According to the theory of Maxwell stress, the electric force on this particle may be calculated from the stress on the above-mentioned surface. For any electric intensity in space there is a corresponding Maxwell stress tensor $\mathbf{T}$ as[3]

$$\mathbf{T} = \frac{1}{4\pi}[\mathbf{EE} - \frac{1}{2}\mathbf{I}(E^2)], \tag{12}$$

where $\mathbf{I}$ is the unit second rank tensor, the elements of $\mathbf{T}$ are

$$T_{ij} = \frac{1}{4\pi}[E_i E_j - \frac{1}{2} I_{ij} E^2], \tag{13}$$

its force element dF_i on an area element ds at the surface is

$$dF_i = \sum_{j=1}^{3} T_{ij} ds_j, \tag{14}$$

where ds_j is the j component of ds. The force F_i on that surface is

$$F_i = \int dF_i. \tag{15}$$

On the spherical surface surrounding the charged particle, the distribution of electric intensity is Eq. (11) with directions along the normal of the surface element ds. Using Eqs. (13), (14) and (15), the total force along z axis on the closed spherical surface is

$$F_z = \oint (T_{zz} ds_z + T_{zx} ds_x + T_{zy} ds_y)$$

$$= \oint \frac{1}{4\pi}\frac{1}{2}(E_z^2 - E_x^2 - E_y^2) ds_z, \qquad (16)$$

since the last two terms integrate to zero owing to their symmetry. Here $ds_z = a^2 \sin\theta_e \cos\theta_e d\theta_e d\phi$ and we may take $\theta \approx \theta_e$ when Z_e is small. Substituting MF of Eq. (6) into Eq. (16) and using $E_z = E_a \cos\theta_e$, $E_x = E_a \sin\theta_e \cos\phi$ and $E_y = E_a \sin\theta_e \sin\phi$, we have

$$\begin{aligned} F_z &= \oint \frac{1}{4\pi}\frac{1}{2}\frac{e^2}{a^4}[1 + \frac{Z_e^2}{(c\Delta t)^2} - 2\frac{Z_e \cos\theta_e}{c\Delta t}]^2 (\cos^2\theta_e - \sin^2\theta_e \cos^2\phi - \sin^2\theta_e \sin^2\phi) a^2 \sin\theta_e \cos\theta_e d\theta_e d\phi \\ &= \oint \frac{e^2}{4a^2}[1 + \frac{Z_e^4}{(c\Delta t)^4} + \frac{4Z_e^2\cos^2\theta_e + 2Z_e^2}{(c\Delta t)^2} - \frac{4Z_e\cos\theta_e}{c\Delta t} - \frac{4Z_e^3\cos\theta_e}{(c\Delta t)^3}](2\cos^2\theta_e - 1)\cos\theta_e \sin\theta_e d\theta_e. \qquad (17) \end{aligned}$$



FIG. 3. The distribution of electric intensities and electric field lines between two spherical surface with radii $r = c\Delta t$ and $r_e = a$.

Here we integrate ϕ from 0 to 2π. The odd terms of $\cos\theta_e$ integrate to zero. Since Z_e is small, the important term of above integration is the first order term of Z_e, those higher order terms can be neglected, thus we have

$$F_z = \oint \frac{e^2}{4a^2}(-\frac{4Z_e\cos\theta_e}{c\Delta t})(2\cos^2\theta_e - 1)\cos\theta_e \sin\theta_e d\theta_e, \qquad (18)$$

thus

$$F_z = -\frac{2e^2}{15a^2}\frac{Z_e}{c\Delta t}. \qquad (19)$$

This displacement dependent F_z is independent of the positive or negative sign of charge e since it's proportional to e^2. This is a kind of displacement dependent restore force induced by the variations of the charge's own electric intensity which follows Coulomb's inverse square law. The magnitude of this force is dependent on the radius a of the spherical surface boundary of the charge, this is different from the stationary electric fields which are independent of the enclosing surface. Here this force is also a temporary force. After the time interval Δt, the charge stops, and as time goes on Δt increase to $\Delta t + t$, where t is the additional time, then Eq. (19) will change into

$$F_z = -\frac{2e^2}{15a^2}\frac{Z_e}{c(\Delta t + t)}. \qquad (20)$$

When t becomes large, F_z reduces to zero.

This problem cannot be treated by usual methods such as the time dependent Coulomb's law[4], the Lienard-Wiechart potential[5] or the Feynman's formula[6] for a charge undergoing an arbitrary translation motion, since the variations of internal structure of the electric fields are not taken into account in these methods. Although the retarded time calculations are used in these methods, the historical conditions of the electric fields are ignored. The problem here is a kind of hereditary electromagnetism[7]. We do not use the integro-differential equation of functional analysis for this problem but the transit effect of displacement is treated in similar way[7].

POSSIBLE EFFECT OF THE DISPLACEMENT DEPENDENT RESTORE FORCE ON THE DYNAMICS OF A FREE ELECTRON

The displacement dependent restore force of Eq. (19) is a kind of self-induced force. For a single free electron with charge e and mass m_e which is not subject to any external force, this self-induced force will affect its dynamical motion. We may take $M_0 r_0$ as the radius a above, where M_0 is an undetermined numerical constant,

r_0 is the classical radius of electron which is defined as[6]

$$r_0 = \frac{e^2}{m_e c^2} \approx 0.82 \times 10^{-13} cm. \tag{21}$$

Substituting $M_0 r_0$ into Eq. (19), we get

$$F_z = -\frac{2e^2}{15(M_0 r_0)^2} \frac{Z_e}{c\Delta t}. \tag{22}$$

For an electron possessing self-sustained harmonic oscillation with angular frequency ω, the time interval Δt may be taken as π/ω, which is half of its period. Then we have

$$F_z = -\frac{2e^2}{15(M_0 r_0)^2} \frac{\omega Z_e}{c\pi} = -\frac{2m_e c\omega Z_e}{15M_0^2 \pi r_0}, \tag{23}$$

since $m_e = e^2/(r_0 c^2)$ according to Eq. (21). If we take $2c/(15M_0^2 \pi r_0) = \omega$, then

$$F_z = -m_e \omega^2 Z_e, \tag{24}$$

which is a standard equation of harmonic oscillation of the free electron. This displacement dependent restore force implies that the electron may have a kind of natural self-oscillation as the "mono-electron oscillator" in a Penning trap of the experiment of Dehmelt et al[8].

The oscillating electron will have oscillating static fields, velocity dependent and acceleration dependent fields, which may store energy and exchange energy between each other[9, 10]. They need not always radiate out their energy through acceleration dependent electromagnetic fields if these fields have standing wave mode solutions[11, 12].

---

[*] lyuqdcn@gmail.com

[1] J. D. Jackson, *Classical Electrodynamics*, 3rd ed. (John Wiley and Sons, 1998) p. 8.

[2] E. H. Williams, J. E. Faller, and H. A. Hill, Phys. Rev. Lett. **26**, 721 (1971).

[3] J. D. Jackson, *Classical Electrodynamics*, 1st ed. (John Wiley and Sons, 1962) p. 193.

[4] D. J. Griffiths and M. A. Heald, Am. J. Phys. **59**, 111 (1991).

[5] W. K. H. Panofsky and M. Philips, *Classical Electricity and Magnetism* (Addison-Wesley, 1962) Chap. 19, p. 341.

[6] R. P. Feynman, R. B. Leighton, and M. Sands, *The Feynman lectures on physics*, Vol. 2 (Addison-Wesley, 1964) Chap. 21.

[7] V. Volterra, *Theory of functionals and of integral and integro-differential equations* (Addison-Wesley, 1959) Chap. VI, p. 194.

[8] D. Wineland, P. Ekstrom, and H. Dehmelt, Phys. Rev. Lett. **31**, 1279 (1973).

[9] J. B. Marion, *Classical Electromagnetic Radiation*, 2nd ed. (Academic Press, 1980) Chap. 8, p. 285.

[10] L. Mandel, J. Opt. Soc. Am. **62**, 1011 (1972).

[11] H. Bateman, *The Mathematical Analysis of Electrical and Optical Wave Motion on the Basis of Maxwell's Equations* (Dover, 1955) Chap. 3, p. 36.

[12] R. B. Adler, L. J. Chu, and R. M. Fano, *Electromagnetic Energy Transmission and Radiation* (John Wiley and Sons, 1960) Chap. 10, p. 566.